

WILDLIFE AND ENVIRONMENTAL PROTECTION

Do not contaminate streams, rivers or waterways by direct application, cleaning of equipment, disposal of wastes and empty container. Product is toxic to animals, bees, fish and other aquatic organisms and may cause long term adverse effects in the aquatic environment.

FIRST AID INSTRUCTIONS

Inhalation: Move the patient to fresh air. Seek medical attention immediately with plenty of water and soap. Seek medical attention if symptoms persist. **Eye Contact:** Wash thoroughly with running water or normal saline solution and obtain medical aid. **It swallowed:** Do not induce vomiting. Do not give anything to unconscious patient. Wash the mouth and keep the patient hydrated. Seek medical attention as soon as possible. **TOXICOLOGICAL INFORMATION/NOT TO PHYSICIAN**

Agrixy 407SL is harmful if swallowed, inhaled or absorbed through the skin and causes moderate eye irritation. **Symptoms of poisoning:** slight eye irritation in case of contact with the eyes or skin irritation. **Antidote:** No specific antidote. **Treat symptomatically and supportively.** **In case of poisoning call toll free number (24hrs) 080072002/0800730030.**

NOTICE TO USERS

This pest control product is to be used only in accordance with the directions on the label. It is an off-label use under the Fast Control Products Act to use or store a pest control product under the directions.

WARRENTY

Seller's guarantee is limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Maelezo

Agrixy 407SL ni kioochoo cha kushikiti 'domey' midew kwa wafidi na 'early and late blight' kwa viazi. **Jinsi inavyoifanya kazi:** Agrixy husafirisha kwa mnea kupitia 'xylem' na phloem 'as huchochua mnea kujikinga dhidi ya kuvu.

JINSI INAYO TUMIKA

CROP	DISEASE	RATES OF APPLICATION
Wardi	Domey/midew	2.5L/ha, 50ml/20L
Viazi	Early and late blight	2.5L/ha, 50ml/20L

Rudia kunyanyaza baada ya siku 7

Kipimo cha maji – Ura 1000

Muda salama wa kukoma – Vazi siku 7

Muda salama wa kuingia paji/ponyuzwa – msaia 12

Kutayarisha mchanganyiko

Weka maji nusu, munoji, ongeza kwanaga kinaochojika cha Agrixy 407SL, huku ukikinga kasha ongeza maji kama mawogendekeza. Hakikishi umrumu mchanganyiko huo siku hyo hyo.

Upaenishi

Agrixy 407SL huchanganika vizuri na madawa ya kuvu wadutu na kuangamiza magonywa isipokuwa ile ya Copper KUZUA UPAENISHI hapaonderwa kwa Agrixy 407SL tumike kwa kuhafidhara na kemikali zilizo na mbinu tofauti za kutanja kazi.



HATARI NA TAHADHARI

Matumizi: Vaa nguo la kujikinga, vatu na vizazi vya mikonzo na uso, unapoyafarisha mchanganyiko wa dawa na maji na unapoyuniza dawa. Jihadiari usipamue muke wa dawa, mchanganyiko wa dawa usitige kwenye macho au kumwagika nguzi na nguzi ulizovalia. Usiinyuzie dawa hii ukielkea upande lipo unatoka. Ustrotie sigara na usile wala kumya wakati unapotuma dawa hii.

Baada ya kazi

Vaa nguzi za kazi na vizazi vya kujikinga, Nawa mikonzo kati ya mapuziko na haru baada ya kazi. Osta bomba uliyotuma kisha Opa na urie mavazi uliyotuma kwa maji na sabuni.

Kuhitaji

Hitari Agrixy 407SL kwenye chombo chake asili kilichofungwa yema na kukwaka, pahali, pakau, pasipo na jito na paanapo hewa vizihi kwenye ghaba laka lililofungwa ambalo watu wassodhiwisha kuingia. Weka mbali na watoto, dhakala, mnyaji na wakaala vya mlingo. Weka mbali na mnyale ya juu, joto jingi na moto.

Ungeji

Suza poochi mara tatu, kitoocha kisha uktupe sehemu ambayo imetengwa na kudhihiwa au kizio mbali na sehemu zilizo na chemu chemu ya maji au ukiregeshe kwenye msambazaji. Msuzo unataa umwagwe mbali na sehemu za maji kuzia machafuko ya mazingira.

Kivujia/mwagika

Vaa mavazi ya kuinga zilizochafuko wa maji au mchanga. Faga kwa kutunga unga wa mbo au mchanga. Kusana kwa kitoo cha mchanga kasha wela ndani ya mikda inayodhihiwa kuondia kemikali. Kwa wingi wa kemikali, tuata maabauri ya msambazaji.

TAHADHARI KWA MAZINGIRA

Agrixy 407SL ni sumu kwa wanyama, nyuki, samaki na vitumbe vya majini. Usichafue vidimbwi au njia za maji. Usiinyuzie kemikali wakati nyuki wanalia kwenye majua. Weka wanyama mbali na sehemu hyo nyunywaza.

MAELEZO YA HUDUMA YA KWANZA

Kwa nguzi: Vaa nguzi na kuinga mara moja kwa maji mengi na sabuni. Kwa macho, Fumbua jicho au macho na kusikilia ghabho cha maji kisha tunda maji mengi ndani ya mado au jicho kwa muda wa dakika 15. Muone daktari. Kivujia: Mpeleleleji, paji/ponyuzwa hawa sari, kwanasumu akivua hewa, mvuso hewa kwa kutuma mdomo wako wa wala. Muone daktari. Kivujia: Mnyaji mwenye kutapika, Kamwe, usimpe kitu chochote inayotuma aliyeveta. Muone daktari. PEEEA, MBERIWA, KIVA, DAKTARI.

MAELEZO KUHUSU SUMU

Agrixy 407SL ni sumu kimerwa, kupunuliwa au kinyagika nguzi na mawalia kilingia machoni.

Dahili za kusunikia

Dahili za sumu ni kama kuwasika kwa nguzi na macho

Makatao

Hakuna makatao maalum

TIBA:

Tibu kulingana na dahili za uponywa

KWA KISA CHA KUSUNIKIA, PIKA SUMU BILA MAIPO MASA 24, 080072002/0800730030

ILANI KWA MTUMIAJI

Dawa hii sarti tumwe kulingana na maagizo yalio kwenye kibandika. Ni hara dhidi ya sheria za modawa ya kuangamiza wadutu wahanichu kutumia au kuhitaji dawa hii katika hali isyo salama. **THIBITISHO** Dhema na yamuzaji ni maelezo yalio kwenye kibandiko ambayo ni lazima yekutawia kiamani. Mounzi lazina akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au maji na lazima sikubali dawa hii katika hali hyo.

GLUE HERE

DRAFT LABEL APPROVED SUBJECT TO
SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 3 MONTHS ONLY

19 JUN 2018

Jhm

PEST CONTROL PRODUCTS BOARD

GLUE HERE

General Information
Agrivyl 407SL is a fungicide with systemic and curative properties for the control of downy mildew in roses and early and late blight in potatoes.

Mode of Action
Agrivyl is a systemic fungicide that is translocated through the xylem and phloem vessels to all parts of the plant. It stimulates plants chemical and physical defenses against disease.

Directions of Application
Agrivyl 407SL should be applied as directed foliar spray. Apply when conditions favor disease development but before the disease occurs and repeat at 7 days interval.

DIRECTION FOR USE

CROP	DISEASE	RATES OF APPLICATION
Roses	Downy mildew	2.5L/ha, 50ml/20L water
Potato	Early and late blight	2.5L/ha, 50ml/20L water

Spray Volume
 Re-entry period
 PH1
 - 1000L
 - 12hrs
 - potatoes 7 days

Preparation of spray mixture
Half fill the tank with water and add the required amount of Agrivyl 407SL mix thoroughly while filling the remainder of the tank with water up to the recommended volume. Ensure that the spray mixture is used on the same day.

Compatibility
Do not apply with copper based fertilizers or fungicides.

Disease resistance
Agrivyl 407SL should be alternated with other products with a different mode of action to prevent development of disease resistant pathogen.

HAZARDS/PRECAUTIONS
Wear protective clothing, overalls, footwear, gloves, and face shield when preparing and using the product. Do not eat, drink or smoke while using it. Avoid inhaling spray mist, contact with skin, eyes and clothing. Do not apply upward and stay out of the spray mist.

After Work
Wash hands before breaks and at the end of work. Change clothes and wash hands and face with soap and water thoroughly. Wash the spraying equipment and any contaminated clothing carefully.

Storage
Keep out of reach of children. Store in tightly closed original container in a cool dry and well ventilated place, under lock and key so that unauthorized persons do not have access. Keep away from food, drinks and animal feeds. Keep away from sunlight, excessive heat and open flame.

Disposal
Triple rinse the pouch and puncture or perforate it. Dispose off in a landfill site approved for pesticides away from water resources or return to stocks/distributor. The residue should be disposed off away from surface water to avoid contamination. Do not re-use the empty container for any other purpose.

DRAFT LABEL APPROVED SUBJECT TO
SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 3 MONTHS ONLY

19 JUN 2018

Jhmj

PEST CONTROL PRODUCTS BOARD

Agrixyl 407SI

FUNGICIDE



A fungicide for the control of early and late blight in potatoes and downy mildew in roses
(Kiua kuu kinachodhibiti baka kwenye viazi na 'downey mildew' kwenye waridi).

COMMERCIAL & AGRICULTURAL CLASS.
(KUNDI LA BIASHARA NA KILIMO.)
GUARANTEE (DHAMANA):
*Mono and di potassium salts of
Phosphorous acid 400g/L + Metalaxyl
70g/L*

**REGISTRATION NO. (NAMBARI YA
USAJILI):** PCPB (CR) 0734
**READ THE LABEL BEFORE USING
(SOMA KIBANDIKO KABLA YA KUTUMIA)**
**KEEP OUT OF REACH OF CHILDREN
(FUNGIA MBALI NA WATOTO)**

Distributor / Wasambazaji:
Amiran Kenya Ltd.
Tel: 0800 720720
Fert@amirankenya.com
www.amirankenya.com

Shelf life: 2 years from date of manufacture when
stored in original unopened container in a cool dry place.
(Maisha Rafuni: Miaka 2 baada ya kutengenezwa kama
imewekwa kwenye chombo chake halisi hakijafunguliwa
mahali pakavu pasipo na joto jingi).


In case of Poisoning call toll free No.
(Kwa kisa cha kusumika piga simu bila
malipo): 0800720021/ 0800730030
(24hrs / masaa 24)

Mft date/Tarehe ya kutengenezwa:
Batch No./Nambari ya Kifurushi:
Expiry date/Tumia kabla ya:

Member Of
Agro Chemical
Association Of Kenya
Registration Number.
AAK. 219



Warning (Ilani)
Very toxic to aquatic life,
may cause long term adverse
effects in the aquatic
environment



PARIJAT **Manufacturer / Watengenezaji:**
PARIJAT INDUSTRIES (INDIA) PVT. L TD.
M - 77, M Block Market, Greater Kailash - II,
New Delhi 110048, India.
Phone: +91 11 41638091/92/93
E-mail product@parijatagrochemicals.com

Net Contents / Ujazo
5 Lts

WHO CLASS II



**HARMFUL/
INADHURU**