accordance with national regulation and legislation.
Pesticide store must be decontaminated by washing with a lot of water.

WILDLIFE AND ENVIRONMENTAL PROTECTION

AZOXY TOP 325SC has low toxicity to mammals, birds and bees. It is highly toxic to fish, algae and other aquatic life and may cause long term adverse effects in the aquatic environment. Do not apply in animal habitats. Keep the product out of aquatic habitats. Do not contaminate irrigation and domestic water supplies by cleaning equipment or waste disposal.

FIRST AID INSTRUCTIONS

If in contact with eye: Hold the eye open and rinse slowly and gently with running water for 15 - 20 minutes.

If in contact with skin: Take off contaminated clothing. Rinse the skin with plenty of water and soap for 15 - 20 minutes.

If inhaled: Move the patient to fresh air.

If swallowed: Rinse the mouth. Do not induce vomiting unless advised by a medical doctor. Do not give anything by mouth to an unconscious person. Seek medical attention immediately.

TOXICOLOGICAL INFORMATION

AZOXY TOP 325SC is harmful when there is repeated exposure at high dose, prolonged contact with skin and with eyes.

Symptoms of Poisoning

AZOXY TOP 325SC may cause mild eye irritation and allergic skin reaction.

Antidote: There is no specific antidote.

Treatment: Treat symptomatically and supportively.

In case of poisoning call toll free telephone no.: 0800720021, 0800730030 (24 hours)

NOTICE TO USERS

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MAELEZO

AZOXY TOP 325SC ni kiua kuvu cha kudhibiti angular leaf spots, anthracnose, na rust kwenye mmea wa French beans, ascochyta leaf spots na powdery mildew kwenye mmea wa snow peas na botrytis na powdery mildew kwenye mmea wa waridi.

INAVYOFANYA KAZI

AZOXY TOP 325SC ni kiua kuvu chenye azoxystrobin na difenoconazole ambazo hufanya kazi pamoja.
Azoxystrobin hufanya kazi kwa njia ya mguzo na kwa kupenya ndani ya matawi ili kuzuia na kuponya kuvu kwa kuitalafiana na ukuwaji. Difenoconazole hufanya kazi kwa kupenya ndani ya matawi na kusambaa kwenye mmea ili kuzuia na kuponya kuvu kwa kuitalafiana na ukuwaji.

MAPENDEKEZO YA MATUMIZI

Mmea	Magonjwa	Kiwango cha Matumizi
French beans	Angular leaf spots, anthracnose, rust	Mililita 500/ Hekta (Mililita 10/Lita 20)
Snow peas	Ascochyta leaf spots, powdery mildew	
Waridi	Botrytis, powdery mildew	Mililita 750 - 1,000/Hekta (Mililita 15 - 20/Lita 20)

MUDA SALAMA WA KUINGIA SHAMBANI: Masaa 6.

MUDA SALAMA WA KUVUNA: French bean, snow peas - Siku 7.

Wakati na Jinsi ya Kutumia

French beans na snow peas: Anza kunyunyiza AZOXY TOP 325SC kabla ya kuona dalili za ugonjwa wakati maua inapotokea ukitumia ratiba ya unyunyizaji wa siku 7 na kiwango cha maji cha lita 1,000 kwa hekta ili kuhakikisha kiua kuvu kimesambazwa vizuri kwenye matawi ya mmea. Usinyunyize AZOXY TOP 325SC zaidi ya mara tatu (3) kwenye mmea bila kubadilisha na viua kuvu vingine.

Waridi: Anza kunyunyiza AZOXY TOP 325SC kabla ya kuona dalili za ugonjwa na wakati hali ya anga inapendelea kuenea kwa magonjwa ukitumia ratiba ya unyunyizaji wa siku 7 na kiwango cha maji cha lita 1,000 kwa hekta ili kuhakikisha kiua kuvu kimesambazwa vizuri kwenve matawi ya mmea. Usinyunyize AZOXY TOP 325SC zaidi ya mara tatu (3) kwenye mmea bila kubadilisha na viua kuvu vingine.

Kutayarisha Kiua Kuvu

Jaza nusu ya bomba la kunyunyiza kwa maji.
Changanya dawa iliyomo kwenye mfuko kabla ya kufungua.
Ongeza kiwango kinachohitajika cha AZOXY TOP 325SC na uchanganye na maji. Ongeza kiasi cha maji kilichosalia na uendelee kukoroga au kutingiza.
Tumia mchanganyiko wa dawa siku uliotengenezwa.

ONYO KWA MTUMIAJI

Utumiaji wa kiua kuvu kimoja au viua kuvu vinavyofanya kazi kwa njia moja mara kwa mara kunaweza kuifanya kiua kuvu kikose nguvu za kukinga mmea dhidi ya magonjwa. Badilisha utumiaji wa AZOXY TOP 325SC na viua kuvu vingine venye njia tofauti ya kuangamiza magonjwa ili kuimarisha utendaji kazi wa kiua kuvu bila kupoteza nguvu za kuangamiza magonjwa.

TAHADHARI

Matumizi

Vaa vizuizi vya mikono na macho, na chombo maalum cha kupumuwa unapotayarisha na ngao la kujikinga, vizuizi vya mikono na macho, chombo maalum cha kupumuwa na viatu unaponyunyiza mchanganyiko wa kiua kuvu na maji. Jihadhari usipumue mvuke wa kiua kuvu, mchanganyiko wa kiua kuvu usiingie kwenye macho au kumwagikia ngozi na nguo ulizovaa. Usinyunyize kiua kuvu hiki ukielekea upande upepo unakotoka. Usivute sigara na usile wala kunywa wakati unapotumia kiua kuvu hiki.

Baada ya Kazi

Vua nguo za kazi na vizuizi vya kujikinga, oga mikono na uso kwa kutumia maji na sabuni. Safisha bomba la kunyunyiza kiua kuvu na mavazi yaliyomwagikiwa na kiua kuvu kwa uangalifu.

Kuhifadhi

Hifadhi kiua kuvu hiki katika chombo chake asili ambacho kimefunikwa sawa sawa na kuwekwa mahali pakavu na pasipo na mwangaza wa jua, joto, moto na mbali na watoto, watu wasioruhusiwa, vyakula, maji na vyakula vya mifugo.

Utupaji

Mkoba ulioisha kiua kuvu ni lazima utobolewe na utupwe kulingana na kanuni na sheria za nchi. Usitumie mkoba wa kiua kuvu kwa njia nyingine. Mchanganyiko wa kiua kuvu ambacho hakijatumika au maji yaliyoosha vyombo ni lazima yatupwe mbali mahali ambapo ni salama, panapokubaliwa na mbali na moto, visima au vidimbwi na chemichemi.

Kiua Kuvu Kikimwagika au Kuvuja

Iwapo kiua kuvu kitamwagika au kuvuja, chukua kiua kuvu kilichomwagika na mchanga ulio na kiua kuvu na uzike mahali ambapo ni salama na mbali na maji. Fungia kiwango kidogo cha kiua kuvu kilichomwagika au mchanga ulio na kiua kuvu mahali ambapo ni salama kabla ya kutupa mbali mahali ambapo pameidhinishwa kulingana na kanuni na sheria za nchi. Ghala la madawa ni lazima lioshwe mara kwa mara ili kutoa takataka ya kiua kuvu kilichomwagika au kuvuja.

MADHARA KWA MAZINGIRA

AZOXY TOP 325SC haina madhara makubwa kwa wanyama, ndege na nyuki. Ni sumu kali kwa zao la mwani, samaki na viumbe vinavyoishi kwenye maji. Usinyunyize kiua kuvu hiki porini. Usichafue mazingara, maji ya kunyunyiza shambani na yale ya kutumia nyumbani kwa kunyunyiza, kumwaga mabaki ya kiua kuvu, kutupa mkoba wa kiua kuvu au kuosha bomba la kunyunyiza kiua kuvu.

MAELEZO YA HUDUMA YA KWANZA

Ikiingia Kwenye macho: Yasafishe macho kwa maji yanayotiririka kwa muda wa dakika 15.

Ikimwagikia ngozi: Vua nguo ulizovaa na uoge mwili kwa kutumia kiasi kikubwa cha maji na sabuni kwa muda wa dakika 15 - 20.

Ikipumuliwa: Pumzisha mjeruhi mahali penye hewa na pasipo na joto nyingi wala baridi.

Ikimezwa: Osha mdomo wa mjeruhi. Usimfanye mjeruhi atapike. Usimpatie mjeruhi ambae amezirahi kitu chochote cha kunywa wala kula. **PELEKA MJERUHI KWA DAKTARI.**

MAELEZO KUHUSU SUMU

AZOXY TOP 325SC ni sumu ikimezwa katika kiwango kikubwa na iwapo itakaa kwenye ngozi au macho kwa muda mrefu.

Dalili za Kusumika

AZOXY TOP 325SC inaweza kusababisha kuwashwa kwa macho na uhisishaji wa ngozi inapokaa kwenye ngozi au macho kwa muda mrefu.

Makata: Hakuna makata maalum.

Maelezo kwa Daktari

Tibu mgonjwa kulingana na dalili za maumivu.

Wakati wa kusumika piga simu bila malipo nambari: 0800720021, 0800730030 (masaa 24) kwa mashauri ya matibabu.

ILANI KWA MTUMIAJI

Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyo salama.

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.

DRAFT LABEL APPROVED SUBJECT TO SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 3 MONTHS ONLY
09 OCT 2018
PEST CONTROL PRODUCTS BOARD

AZOXY TOP 325SC

Fungicide for control of angular leaf spots, anthracnose and rust in French beans, ascochyta leaf spots and powdery mildew in snow peas and botrytis and powdery mildew in roses

(Kiua kuvu ya 'angular leaf spots', 'anthracnose' na 'rust' kwenye mimea ya French beans, 'ascochyta leaf spots' na 'powdery mildew' kwenye mimea ya snow peas na 'botrytis' na 'powdery mildew' kwenye mimea ya waridi)

COMMERCIAL AND AGRICULTURAL CLASS
(KUNDI LA BIASHARA NA KILIMO)

REG. NO: (NAMBARI YA USAJILI): PCPB (CR) 1389

GUARANTEE: Azoxystrobin ......200 g/L
(DHAMANA) Difenoconazole ...125 g/L

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USE

(WEKA MBALI NA WATOTO)
(SOMA KIBANDIKO KABLA YA KUTUMIA)

Shelf life: 2 years from date of manufacture when stored in original unopened container in a cool dry place.

(**Maisha rafuni:** Miaka 2 baada ya kutengenezwa ikihifadhiwa kwenye chombo chake halisi kisichofunguliwa, mahali pakavu pasipo na joto nyingi)

Batch No.: ....
(Nambari ya kifurusi)

Date of Manufacture: ....
(Tarehe ya Kutengenezwa)

Expiry Date: ....
(Tumia kabla ya)

INCASE OF POISONING CALL TOLL FREE TELEPHONE NO.:
0800720021, 0800730030 (24 HOURS)
(WAKATI WA KUSUMIKA PIGA SIMU BILA MALIPO NAMBARI (MASAA 24) KWA MASHAURI YA MATIBABU)

Content (Ujazo): 500 ml

Agent (Wasambazaji):
AGROSOLUTIONS LTD.
Off Mombasa Road
P.O. BOX 53418 - 00200 Nairobi
Tel: (020) 2043358, 0724625315,
Email:info@agrosolutions.co.ke

Manufacturer (Watengenezaji):
ZHEJIANG BOSST CROPSCIENCE CO., LTD.
NO.1 Fangjiadai Road,
Haiyan Economic
Dev. Zone, Haiyan,
Zhejiang, China

HARMFUL (INADHURU)

DESCRIPTION

AZOXY TOP 325SC is a broad-spectrum fungicide for control of angular leaf spots, anthracnose and rust in French beans, ascochyta leaf spots and powdery mildew in snow peas and botrytis and powdery mildew in roses.

MODE OF ACTION

AZOXY TOP 325SC contains azoxystrobin and difenoconazole that complement each other.

Azoxystrobin has preventive, curative and eradicant activity with contact and translaminar properties and acts by inhibiting respiration in the mitochondrial in the fungi, thereby inhibiting spore germination, germ tube development and sporulation.

Difenoconazole has translaminar and systemic action with preventive and curative properties and acts on fungi during penetration and haustoria formation. It stops development of fungi by interfering with the biosynthesis of sterols in cell membranes.

DIRECTION FOR USE

Crop	Disease	Application Rate
French beans	Angular leaf spots, anthracnose, rust	500 ml/Ha (10 ml/20L)
Snow peas	Ascochyta leaf spot, powdery mildew,	
Roses	Botrytis, powdery mildew	750 - 1000 ml/Ha (15 - 20 m1/20L)

PRE-HARVEST INTERVAL

French beans, snow peas - 7 days.

RE-ENTRY PERIOD: 6 hours.

Application Timing and Technique

French beans and snow peas: Start spray application before disease condition appears at flower initiation stage at 7 days interval and with adequate spray volume of 1,000 L of water/Ha to ensure thorough coverage of plant surface. Do not apply more than three (3) successive sprays in a season.

Roses: Start spray application before disease condition appears or when weather condition favour disease development at 7 days interval. Apply with adequate spray volume of 1,000 L of water/Ha to ensure thorough coverage of plant surface.

Preparation of Spray Mixture

Half fill the spray tank with water. Roll or shake the AZOXY TOP 325SC container before use and add the required amount in the spray tank. Fill the spray tank with water and stir or agitate to uniformity.

Use the spray mixture the same day of preparation.

DISEASE RESISTANCE

Continuous use of AZOXY TOP 325SC or in alternation with fungicides of similar mode of action may predispose the target fungi to develop resistance to AZOXY TOP 325SC or other fungicides with similar mode of action. Alternate treatments of AZOXY TOP 325SC with fungicides of different mode of action to delay the development of disease resistance.

HAZARDS/PRECAUTIONS

Handling

Wear protective hand gloves, face-shield, overall and impermeable boots when preparing and applying the spray mixture. Avoid breathing the spray mist, contact with skin, eyes and clothing. Do not apply upwind and stay out of spray mist. Do not smoke, drink or eat while handling the product.

After Work

Change clothes and wash hands and face thoroughly with soap and water. Wash the spray equipment and any contaminated clothing carefully.

Storage

Store in original tightly closed (or sealed) container, in a cool dry, well ventilated place, and out of reach of children, unauthorized persons and animals, away from food, drinks and animal feed and fire or heat sources.

Disposal

Triple rinse, puncture or crush empty containers before disposal in a safe and authorized place in accordance with national regulations and legislation. Do not re-use empty containers for any other purpose. Pesticide spray mixture or rinsate that cannot be used must be disposed off where it cannot contaminate surface water and ditches.

Pesticide Leakage and Spillage

In the event of spillage, collect the product and contaminated soil layer and bury in an authorized place away from water supplies. For small spillage, collect in absorbent material. The contaminated absorbent material must be stored in marked waste bins prior to disposal off in an authorized place away from water supplies in

DRAFT LABEL APPROVED
SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 6 MONTHS ONLY
09 OCT 2018
PEST CONTROL PRODUCTS BOARD