General information: DIFFERENCE 250EC is a systemic fungicide with protective, curative and eradicant properties recommended for the control of powdery mildew & botrytis on Roses.
Mode of action DIFFERENCE 250EC is taken up by the plant and acts on the fungal pathogen during penetration and haustoria formation. It stops development of the fungi by interfering with the biosynthesis of sterols in cell membranes. DIFFERENCE 250 EC is a sterol demethylation inhibitor.
Direction for use

CROP	DISEASE	APPLICATION RATE	SPRAY INTERVAL	R.E.I
Roses	Powdery mildew	0.375L/Ha/1000L water, 37.5ml/100L water, 7.5ml/20L water	7-14 days	6 hrs/after spray dies
	Botrytis	0.5-0.625L/Ha/1000L water, 50-62.5ml/100L water, 10ml/20L water	10-14 days	

Preparation of Spray Mixture Add the required amount of DIFFERENCE 250EC to half filled spray tank and agitate while adding the remaining water. Maintain good agitation throughout the spraying operation. Do not allow the spray mixture to stand overnight or for long periods.
Application timing It is recommended to apply DIFFERENCE 250EC early enough to avoid irreversible crop damage and build up of the disease. A maximum of 3 applications of DIFFERENCE 250EC is recommended from onset of disease.
Applications Technique DIFFERENCE 250EC can be applied with all types of ground spray equipments and should use sufficient water volume (1000L/Ha) for adequate coverage and canopy penetration.

HAZARDS/ PRECAUTIONS
Wear protective clothing namely overalls, footwear, gloves, and face shield when preparing and using the product. Do not eat, drink or smoke when working with the product.
After work Wash hands before breaks and at end of work. Change clothes and wash hands and face with soap and water thoroughly. Wash the spraying equipment and any contaminated clothing carefully.
Storage Keep out of reach of children. Store in tightly closed original container in a cool,dry and well ventilated place, under lock and key so that unauthorized persons do not have access. Keep away from food, drinks and animal feeding stuffs.
Disposal Triple rinse, puncture or crush empty containers before disposal in a safe and authorized place in accordance with National regulations on chemical disposal. Do not re-use empty containers for any other purpose. Pesticide spray mixture or rinsate that cannot be used must be disposed where it cannot contaminate surface water and ditches.
Pesticide Leakage and Spillage In event of heavy spillage, collect the contaminated soil layer and dispose in an authorized place away from water supplies. Soaking in sand, soil or sawdust may collect small spillage. Contaminated absorbent material must be stored in marked waste bins prior to disposal in authorized landfill site. The pesticide store must be decontaminated by washing with a lot of water.
Wildlife and Environmental Protection
DIFFERENCE 250EC is toxic to fish, aquatic life and wildlife. Do not apply directly to water when disposing or to areas of runoff or contaminate water when disposing of equipment wash water or rinsate.

FIRST AID INSTRUCTIONS
Skin contact: Remove affected clothing and wash all exposed skin area with soap and water, followed by warm water rinse. Consult a doctor in the event of any complaints.
Eye contact: Wash off with plenty of water (15 min.) If pain persists, consult an eye specialist.
If inhaled: Remove victim to fresh air. Consult a doctor in the event of any complaints.
If swallowed: Do not induce vomiting. Wash out mouth with plenty of water. Never give anything by mouth to an unconscious person. Consult a doctor in the event of any complaints.

TOXICOLOGICAL INFORMATION
Symptoms of Poisoning: Headache, dizziness, slight skin & eye irritant, weakness, abdominal pain & nausea. **Antidote:** There is no specific antidote.
NOTE TO PHYSICIAN: Treat symptomatically.
In case of Poisoning call toll free number: 0800720021/0800730030 (24hrs)

NOTICE TO USERS
This pest control product is to be used only in accordance with the directions on the label.
It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.
WARRANTY Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Amiran

Difference

250EC

FUNGICIDE

A fungicide for Control of Powdery Mildew & Botrytis on Roses
(Kiua kuvu kinacho dhibiti kuvu ya unga na Botraiti kwenye waridi).

COMMERCIAL & AGRICULTURAL CLASS.
(KUNDI LA KILIMO NA BIASHARA.)
GUARANTEE (DHAMANA): *Difenoconazole 250g/L*
REGISTRATION NO. (NAMBARI YA USAJILI):
PCPB (CR) 1249

READ THE LABEL BEFORE USING
(SOMA KIBANDIKO KABLA YA KUTUMIA)

KEEP OUT OF REACH OF CHILDREN
(WEKA MBALI NA WATOTO)

Shelf life: 2 years from date of manufacture when stored in original unopened containers in a cool dry place.
(Maisha Rafuni: Miaka 2 tokea ya kutengenezwa kama imewekwa kwenye chombo chake halisi hakijafunguliwa mahali pakavu pasipo na joto jingi).

Mft date/Tarehe ya kutengenezwa:
Batch No./Nambari ya Kifurushi:
Expiry date/Tumia kabla ya:

In case of Poisoning call toll free No.
(Kwa kisa cha kusumika piga simu bila malipo-masaa 24):
0800720021/ 0800730030(24hrs)

Member Of Agro Chemical Association Of Kenya Registration Number. AAK 219

aak

Distributed By / Wasambazaji:
Amiran Kenya Ltd.
Tel: 0800 720720
Fert@amirankenya.com
www.amirankenya.com

Arysta

Manufacturer / Watengenezaji:
Arysta LifeScience BELGIUM.

Net Contents / Ujazo **1 Lt**

WHO CLASS II HARMFUL (HADHURU)

Maelezo DIFFERENCE 250 EC ni kiua kuvu kinacho zuia, kuangamiza na kudhibiti kuvu ya unga na Botraiti kwenye waridi.
Jinsi Inavyofanya kazi DIFFERENCE 250EC hupenya ndani ya mmea na kudhibiti ukungu wakati zinapoceneya kwa mmea. Inazuia kungu kwa kuharibu chembe za uhai.

MAPENDEKEZO YA MATUMIZI

MIMEA	WADUDU	VIPIMO VYA MATUMIZI	WAKATI WA KURUDIA	SUBIRI
Waridi	Kuvu ya unga	Lita 0.375/Hekta, lita 1000 za maji, mililita 37.5 kwa lita 100 za maji, mililita 7.5/lita 20 za maji Lita	Siku 7-14	Masaa 6/ Baada ya dawa kukauka
	Botraiti	0.5-0.625/Hekta, lita 1000 za maji, mililita 50-62.5 kwa lita 100 za maji, mililita 10-12/lita 20 za maji	Siku 10-14 days	

Kutayarisha Dawa Pima kiasi cha DIFFERENCE 250EC kwa bomba lililojazwa kiasi cha maji huku ukikoroga. Kisha jaza Kiwango cha maji kinachohitajika ilikuwa na mchanganyiko unaofaa. Endelea kutikisa bomba unaponyunyizia. Tumia mchanganyiko wa dawa siku hiyo ulioitayarisha.
Wakati wa Kutumia Inapendekezwa kutumia dawa hii mapema kuzuia uharibifu wa mmea na kuenea kwa ugonjwa. Pia inapendekezwa kutumia Difference 250EC mara 3 baada ya ugonjwa kudhihirisha.
Jinsi ya Kutumia DIFFERENCE 250EC yaweza kunyunyizia kupitia aina nyingi za bomba na inahimizwa kutumia juzuu la maji (lita 1000 kwa hekta) ili mchanganyiko was dawa kutosha na kupenya kwa mmea.
HATARI NA TAHADHARI
Vaa nguo za kujikinga, viatu na vizuizi vya mikono na osho (face shield), unapotayarisha mchanganyiko wa dawa na maji na unaponyunyiza dawa. Jihadhari usipumue mvuke wa dawa, mchanganyiko wa dawa usingie kwenye macho au kumwagikia ngozi na nguo ulizovalia. Usinyunyizie dawa hii ukielekeza upande upepo unatoka. Usivute sigara na usile wala kunywa wakati unapotumia dawa hii.
Baada ya kazi: Vua nguo za kazi na vizuizi vya kujikinga, oga mikono na uso kwa kutumia sabuni na maji. Safisha bomba la kunyunyizia dawa na mavasi yaliyomwagikiwa na dawa.
Kuhifadhi Hifadhi dawa hii katika chombo chake asili ambacho kimefunikwa na kuwekwa kwenye ghala la madawa lililoidhinishwa. Weka mbali na watoto, chakula, vinywaji na vyakula vya mifugo.
Utupaji Safisha mkoba wa dawa mara tatu kisha utoboe au ubonyeze na utupilie mbali na maji ambayo ni salama na panapoidhinishwa kulingana na sheria za utupaji. Mchanganyiko wa dawa ambao hautumiki au maji yaliyoosha vyombo ni lazima yatupiliwe mahali salama na ambapo panaidhinishwa. Dawa ambayo haiwezi kutumiwa ni lazima itupiliwe mbali mahali ambapo ni salama, panapokubaliwa, na mbali na maji. Usitumie mkoba wa dawa kwa matumizi mengine yoyote.
Dawa Ikimwagika au Kuvuja Chukua mchanga uliomwagikiwa na dawa, utupe mahali ambapo ni salama na mbali na maji. Dawa ikivunja, iloweshe kwenye sehemu ndogo ya mchanga au 'sawdust' kisha ufungie mahali ambapo ni salama kabla ya kutupilia mbali mahali ambapo panaidhinishwa. Ghala la madawa ni lazima lioshwe mara kwa mara ili kutoa takataka ya dawa iliomwagika au kuvunja.
MADHARA KWA MAZINGIRA
DIFFERENCE 250EC inadhuru samaki na viumbe vingine vya majini na wanyama poli. Usichafue visima au mito kwa kunyunyizia, kuosha vyombo, kumwaga mabaki au mkoba ya dawa hii.

MAELEZO YA HUDUMA YA KWANZA
Ikipumuliwa: Toa mgonjwa mahali penye hatari na umpumzishe mahali ambapo ni salama. Kama mjeruhi anashida ya kupumua, msaidie apumue, kisha umpeleke kwa daktari.
Ikimwagikia ngozi: Vua mavazi ya kujikinga na uoge kwa kutumia maji safi na sabuni. Iwapo maumivu itaendelea, muone daktari.
Ikingia kwa macho: Fungua macho na uyasafishe kwa maji safi yanayotiririka kwa muda wa dakika 15. Iwapo maumivu itaendelea, muone daktari. **Ikimezwa:** Usimfanye mgonjwa atiye katika hali maututi kutapike. Muoshe mdomo, kisha umuone daktari. PELEKA MJERUHIWA KWA DAKTARI
MAELEZO KWA DAKTARI: Tibu mgonjwa kulingana na dalili za maumivu.
Kwa kisa cha kusumika piga simu bila malipo (masaa 24): 0800720021/0800730030
ILANI KWA MTUMIAJI
Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyo salama.
THIBITISHO Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.