

**READ LABEL BEFORE USE(SOMA KIBANDIKO KABLA YA KUTUMIA)
KEEP LOCKED OUT OF REACH OF CHILDREN(FUNGIA MBALI NA WATOTO)**



Fungicide for the control of Botrytis in Roses: Early blight in tomatoes, soft rot and black leg on broccoli, angular leaf spot in French beans and runner beans (Kiua kuvu cha kuangamizaugoniwa wa 'Botrytis' kwenye waridi, 'Early blight' kwenye nyanya, 'soft rot' na 'black leg' kwenye 'broccoli' na 'angular leaf spot' kwenye michiri na maharagwe)

GUARANTEE (DHAMANA): Carbendazim 250g/L & Pyrimethanil 50g/L

Rimeta Gold 300SC is a Trademark of (alama ya biashara ya): Greenlife Crop Protection Africa Ltd.
**COMMERCIAL & AGRICULTURAL CLASS (KUNDI LA BIAHARA NA KILIMO)
REGISTRATION NO. (NAMBARI YA USAJILI): PCPB (CR) 24/0001/0800730030**

FUNGICIDE

**In case of poisoning, call (24hr) toll free number)
Wakati wa kusumika, piga simu(masa 24)
0800720021/0800730030**

**DRUGS BOARD APPROVED SUBJECT TO
SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 3 MONTHS ONLY
27 APR 2018
PEST CONTROL PRODUCTS BOARD**

Manufactured by (Watengenezaji):
Yantai Keda Chemical Co. Ltd. Zhaoyuan, Shandong, China



Greenlife
CROP PROTECTION AFRICA
Net Contents
(Kiasi) **1L**

Distributed by
(Wasambazaji):



#.361

Batch No. _____
(Nambani ya kifurushi)
Date of Manufacture (Tarehe ya kutengenezwa): _____
Expiry Date. (Tumia Kabla ya): _____

Shelf Life (Maisha Rafani): 2 years from the date of manufacture when stored in original unopened container in a cool dry place.
(Mfaka miwili baada ya tarehe ya kutengenezwa kama imehifadhiwa kwenye chombo chake asili kusichofunguliwa pahali pakavu pasipo na joto jingi)



CAUTION
TAHADHARI



DESCRIPTION: Fungicide for the control of Botrytis in Roses; Early blight in tomatoes; soft rot and black leg on broccoli; angular leaf spot in French beans and runner beans
MODE OF ACTION: Acts by inhibiting the infection process; germ tube elongation, mycelium growth and sporulation as well as inhibiting fungal mitotic microtubule formation. Carbendazim belongs to Benzimidazole class while Pyrimethanil belong to Anilino-pyrimidine class.

APPLICATION RATES

Crop	Disease	Rate
Roses	Botrytis	1.0 - 2.0L/ha in 1000 L of water
Frenchbeans	Angular leaf spot	(20 - 40 ml in 20 L of water)
Runner beans	Angular leaf spot	
Tomatoes	Early blight	
Broccoli	Soft rot, Black leg	

PHI: French beans and runner beans 14 days, Tomatoes and Broccoli 7 days.

APPLICATION METHOD: Apply RIMETA GOLD 300SC to the point of runoff using 1,000 litres of water per hectare to ensure adequate penetration and coverage of the crop foliage and flowers.

APPLICATION TIMING: Start applying RIMETA GOLD 300SC when weather conditions favour disease development or when disease symptoms are noted.

PREPARATION OF SPRAY MIXTURE: Fill spray tank with half the amount of required water. Add required amount of RIMETA GOLD 300SC and fill the water to the required level. Agitate to ensure thorough mixing. Use spray mixture on the same day of preparation.

RE-ENTRY PERIOD: Do not enter treated areas for 6 hours.

HAZARD/PRECAUTION SAFETY PRECAUTIONS:

Wear overall, impermeable gloves, boots, and face shield when mixing and spraying and when cleaning application equipment. Avoid inhaling or contact of product with skin, eyes and clothing. Do not apply upwind. Do not smoke, drink or eat when handling this product. After work, carefully clean the spraying equipment, remove contaminated clothing and wash hands and face thoroughly with water and soap.

STORAGE: Store in original container in a dry storage area. Keep container tightly closed. Keep away from food, drink and animal feed. Keep out of reach of children.

DISPOSAL: Triple rinse and puncture empty containers and dispose in a safe designated place away from water sources in accordance with National Regulations on disposal. Do not reuse empty containers for any other purposes.

ENVIRONMENTAL HAZARDS: RIMETA GOLD 300SC is toxic to fish and other aquatic organisms. Do not contaminate streams, rivers or waterways with the chemical or used containers. Do not spray while bees are actively foraging.

FIRST AID INSTRUCTIONS

Inhalation: Move victim to fresh air. If victim is not breathing, give artificial respiration.

Skin contact: Wash off immediately with soap and plenty of water.

Eye contact: Immediately rinse eyes with plenty of water for 15-20 minutes.

Ingestion: Wash the mouth with plenty of water. Do not induce vomiting. Keep patient at rest.

TOXICOLOGICAL INFORMATION: Harmful if swallowed, inhaled or in contact with the skin.
Symptoms of poisoning: Throat and eye irritation.
Antidote: No known antidote.
Note to the Physician: Treatment should be symptomatic.

In case of poisoning, call (24hr) toll free number 0800720021 or 0800730030

NOTICE TO USERS: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MAELEZO: Dawa ya kuangamiza ugonjwa wa Botrytis kwenye waridi, Early blight kwenye nyanya, Soft rot na black leg kwenye broccoli na angular leaf spot kwenye michiri na maharagwe
MAVOFANUA KAZI: Huzua kuanza, kukua na kuenea kwa viini vinyasomabaza magonywa, Carbendazim ni ya kundi la Benzimidazole. Ithali Pyrimethanil ni ya kundi la Anilino-pyrimidine.

KIPIMO CHA KUTUMIA

Mimea	Magonjwa	Kipimo
Waridi	Botrytis	Lita 1.0 - 2.0 kwa hektari kwa lita 1000 za maji
Michiri	Angular leaf spot	(milita 20 - 40 kwa lita 20 za maji)
Maharagwe	Angular leaf spot	
Nyanya	Early blight	
Broccoli	Soft rot, Black leg	

MUDA SALAMA KABLA YA KUVUNA: Michiri na maharagwe, siku 14. Nyanya na Broccoli siku 7.

JINSI YA KUTUMIA: Nyunyiza mimea kwa kutumia kiasi cha lita 1000 za mchanganyiko wa dawa na maji kwa hektari moja, ili kuhakikisha dawa imesambazwa vizuri kwenye matawi na mawa.

WAKATI WA KUTUMIA: Anza kutumia RIMETA GOLD 300SC wakati dalili za magonywa zinapoonakana au wakati hali ya anga inapendelea kuokea kwa magonywa.

JINSI YA KUCHANGANWA: Ila nusu ya kiasi cha maji yanayohitajika kwenye bomba halafu ongeza kivango cha RIMETA GOLD 300SC kinachohitajika. Kisha ongeza maji yaliyobakia huku ukikoroga vizuri.

TAHADHARI: Yaa mavazi ya kinga ya uso, miitiko ya raba ya mikonu, ovaroli, na vitu visivyopenyeka dawa unapochanganya na kunyunyiza dawa hii. Usinyunyize dawa hii ukiteleka unakotoka urepo. Eruka dawa kuingia kwenye macho, mapua au kugusana na ngozi. Usile, kunywa wala kuvuta sigara unapoitumia dawa hii. Baada ya kutumia dawa, osha vizuri bomba ya kunyunyiza dawa na mavazi ya kujikinga. Oga mikono na uso kwa maji na sabuni.

KUHIFADHI: Dawa hii yafaa kuhifadhiwa kwenye chombo chake asili, pahali pakavu, pasipo na joto jingi huku ukipenka kuchanganya na vyakula. Weka mbali na watoto.

UTUPAJI: Chombo kitupu cha dawa hii chafaa kutobolewa na kutupwa kulingana na sheria na kanuni za utupaji taka pahali palipoingewa, mbali na chemchemi au hiradhi za maji. Usitumie chombo kitupu kwa kazi zingine zozote.

ADHARI KWA MAZINGIRA: RIMETA GOLD 300SC, ya kudhuru samaki na viumbe wa maji. Kwa hivyo dawa yenyewe, mabaki yake au chombo kitupu hazijai kutimwa au kutupwa kwa njia tekeleyochafua visima, mita, vidimbiwi, au chemchemi za maji. Usitumie dawa hii wakati nyuki wanapo onekana kwa wingi shambani

HUDUMA YA KWANZA:

Ikwitwa kwa mapanuzi: Eweka mihadhiwa palipo na hewa safi na umtuzi. Kiwa hapurui, masidie kupumua.

Ikgusana na ngozi: Safisha ngozi vizuri kwa maji mengi na sabuni.

Ikingia kwa macho: Fungua na kugaocha macho na maji mengi kwa dakika 15.

Ikimezwa: Osha mdomo kwa maji. Usimfanye kutapika. Mutuze mihadhiwa. **PELEKA MWERUHI KWA DAKTARI**

MAELEZO KUHUSU SUMU: Inadhiru ikimezwa, ikivutwa kwa mapua au kigusa ngozi.

Dalili za kusumikar: kuwashwa kwa koo na macho.

Makata: Hakuna tiba maalum.

Aria kwa daktari: Thiu kulingana na dalili za kusumika.

Wakati wa dhuru, piga simu (masasa 24) bila malipo kwa 0800720021/0800730030

ILANI KWA MTUMIAJI: Dawa hii sharti itumwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za maddawa ya kuangamiza wadudu waharibi kutumia au kuhifadhi dawa hii katika hali isiyosajama.

THIBITISHO: Dhamana ya mruzaji ni maelezo yaliyo kwenye kibandiko ambayo lazima yafutwe kikamilifu. Mhunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au maji na lazima akubali dawa katika hali hivyo



Greenlife
CROP PROTECTION AFRICA
P. O. Box 2092 - 00100, Nairobi, Kenya. Fax: +254 738 980287
Mtc: +254 735 54544 +254 722 56558. Tel: +254 20 244220/2
info@greenlife.co.ke | www.greenlife.co.ke

Your Growth. Our Growth

Net Contents
(Kiasi) **1L**