

**READ LABEL BEFORE USE/SOMA KIBANDIKO KABLA YA KUTUMIA)**  
**KEEP LOCKED OUT OF REACH OF CHILDREN/FUNGIA MBALI NA WATOTO)**



Insecticide Seed dresser for the control of Russian wheat aphid and Green bug in Wheat; and early pests of Cotton (aphids, whiteflies, mealybugs and jassids); Stalk borer, cutworms, thrips, beetles & aphids in seed maize (Kua dudu kinachotibu mbevu na kuzuia wadudu waharibifu kama vile 'Russian wheat aphid' na 'Green bug' kwenye ngano na wadudu wa mapema kwenye pamba, 'aphids', 'whiteflies', 'mealybugs' na 'jassids'); 'Stalk borer', 'cutworms', 'thrips', 'beetles' na 'aphids' kwenye mbevu za mahindi

**GUARANTEE (DHAMANA): Imidacloprid 600 g/L**

Shield 600 FS is a Trademark of (alama ya biashara ya): Greenlife Crop Protection Africa.

COMMERCIAL & AGRICULTURAL CLASS (KUNDI LA BIASHARA NA KILIMO)  
 REGISTRATION NO. (NAMBARI YA USAJILI): PCPB (CR)1241

## INSECTICIDE

In case of poisoning call (24hr) toll free number  
 Wakati wa kusumika, piga simu(masa 24) bila malipo kwa namba 0800720021/0800730030

27 JUL 2018

DRAFT LABEL APPROVED SUBJECT TO THE SUBMISSION OF FINAL LABEL AND PACKAGE DESIGN. VALID FOR 3 MONTHS ONLY

Distributed by  
 (Wasambazaji):



#.361



**Greenlife**  
 CROP PROTECTION AFRICA  
 Net Contents  
 (Kiasi) 1L

Batch No. \_\_\_\_\_  
 (Namba ya kifunushi)  
 Date of Manufacture (Tarehe ya kutengenezwa): \_\_\_\_\_  
 Expiry Date, (Tumia Kabla ya): \_\_\_\_\_

Shelf life (Maisha Rafuni): 2 years from the date of manufacture when stored in original unopened container in a cool dry place. (Mlaka miwili baada ya tarehe ya kutengenezwa kama imehifadhiwa kwenye chombo chake asili kusichofunguliwa pahali pakavu pasipo na joto jingi)

**CAUTION**  
 (Tahadhari)



**DESCRIPTION:** SHIELD® 600 FS is a seed dressing insecticide with residual effect for control of Russian wheat aphid and Green bug in Wheat; and early pests of Cotton (aphids, whiteflies, mealybugs and jassids); Stalk borer, cutworms, thrips, beetles & aphids in seed maize.  
**MODE OF ACTION:** SHIELD® 600 FS has contact and stomach poison action. It interferes with the nerve transmission system of insects. It belongs to neonicotinoid class.

**APPLICATION RATES**

Crops	Pest	Rate
Maize	Stalk borer, cutworms, thrips, beetles & aphids	3 ml/Kg seeds mixed with 20 ml of water
Wheat	Russian wheat aphid & Green bug	200ml/1.6L of water/ 100kg of seeds Apply once as seed dresser and dry under shade for one hour
Cotton	Aphids, whiteflies & mealybugs	2ml/1Kg of seeds Apply once as seed dresser and dry under shade for one hour
	Jassids	2.5ml/1Kg of seeds Apply once as seed dresser and dry under shade for one hour

**PRE-HARVEST INTERVAL:** Not required when used as directed.  
**RE-ENTRY PERIOD:** Not required when used as directed.

**APPLICATION TECHNIQUE:** After adding the required quantity of SHIELD® 600 FS mixed with the required volume of water, close the lid and rotate the drum slowly for 3-5 minutes to ensure that each seed is properly covered with the seed dressing and allow to dry for 24 hours before planting.

**HAZARD/PRECAUTION**

**SAFETY PRECAUTIONS:** Avoid contact with skin and eyes. When using, do not eat, drink or smoke. Wear overalls, safety glasses and impervious gloves. Wash contaminated clothing after protective equipment before storage or re-use.

**STORAGE:** Store in the closed, original container in a cool, well-ventilated area. Do not mix with foodstuff or feed in storage. Keep out of reach of children.

**DISPOSAL:** Triple rinse the container then puncture empty containers and dispose in a safe designated place away from water sources and in accordance with national regulation and legislation on disposal. Do not reuse empty containers for any other purposes.

**ENVIRONMENTAL HAZARDS:** SHIELD® 600 FS is toxic to fish and wildlife. Do not contaminate streams, rivers or waterways with the chemical or used containers. Treated seed must be planted properly and totally covered with soil. No seed should be left in places where it can be eaten by birds.

**FIRST AID TREATMENT:**

**Inhalation:** If inhaled, move to fresh air and keep at rest. If not breathing, give artificial respiration.

**Skin contact:** Wash off immediately with running water for 20 minutes.

**Eye contact:** Immediately rinse eyes with plenty of water for 15 minutes.

**Ingestion:** Wash out mouth with water. Do not induce vomiting. Keep patient at rest.

**TOXICOLOGICAL INFORMATION:**

**Symptoms of poisoning:** Irritation of eyes, skin and mucous membranes allergy skin rashes at prolonged contact.

**Antidote:** No known antidote. Seek medical advice.

**Note to the physician:** Treatment should be symptomatic.

**In case of poisoning, call (24hr) toll free number 0800720021 or 0800730030**

**NOTICE TO USERS:** This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

**WARRANTY:** Seller's guarantee is limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

**MAELEZO:** SHIELD® 600 FS ni dawa inayotibu mbevu na kuzuia wadudu waharibifu kama 'Russian wheat aphid' na 'Green bug' kwenye ngano na wadudu wa mapema kwenye pamba, 'aphids', 'whiteflies', 'mealybugs' na 'jassids'; 'Stalk borer', 'cutworms', 'thrips', 'beetles' na 'aphids' kwenye mbevu za mahindi.  
**INAVOFANYA KAZI:** SHIELD® 600 FS hua wadudu kwa kupenya ndani ya mmea kutoka kwenye matawi hadi kwenye mizizi. Ni ya kundi la neonicotinoid.

**KIPIMO CHA KUTUMIA**

Mmea	Wadudu	Kipimo
Mahindi	Stalk borer, 'cutworms', 'thrips', 'beetles' na 'aphids'	Millilita 3 kwa kilo 1 ya mbevu lichoanganywa na mililita 20 ya maji
Ngano	'Russian wheat aphid' na 'Green bug'	Millilita 200 kwa lita 1.6 za maji kwa kilo 100 za mbevu (Nyunyuzia mara moja kisha uzache mbevu kwa kiwili kwa mda wa saa 1)
Pamba	'Aphids', 'whiteflies' na 'Mealybugs'	Millilita 2 kwa kilo 1 za mbevu (Nyunyuzia mara moja kisha uzache mbevu kwa kiwili kwa mda wa saa 1)
	'Jassids'	Millilita 2.5 kwa kilo 1 za mbevu (Nyunyuzia mara moja kisha uzache mbevu kwa kiwili kwa mda wa saa 1)

**MUDA SALAMA KABLA KUVUNA:** Hawatakiwi wakati kutumika kama ilivyoagizwa.

**MUDA SALAMA WA KUNINGIA PALIPONYUNYIZIWA:** Hawatakiwi wakati kutumika kama ilivyoagizwa.

**JINSI YA KUCHANGANYWA:** Pima kiasi cha dawa kinachotijika. Changanya na kipimo halisi cha maji. Weka mbevu na uchanganye vizuri kwa muda wa dakika 3-5. Toa mbevu na ukaushe kwa muda wa masaa 24 kabla ya kupanda.

**TAHADHARI:** Euka dawa kuingia kwenye macho, au kugusana na ngozi. Usile, kunywa wala kuvuta sigara unapoitumia dawa hii. Yaa mavazi ya kinga k.v. kinga ya uso, mifuko ya raba ya mlkono, na overall unapochanganya na kunyunyizia dawa hii. Baada ya kutumia dawa, osha vizuri mavazi ya kujikinga.

**KUHIFADHI:** Dawa hii yafaa kuhifadhiwa kwenye chombo chake asili, pahali pakavu. Usihifadhi pantoga na yakula vya watu au mitugo. Weka mbali na watoto.

**UTUPAJI:** Chombo kitupu cha dawa hii chafaa kusafishwa mara tatu, kutobolewa na kutupwa pahali palipotengwa, mbali na chemchemi au hifadhi za maji kulingana na kanuni na sheria za nchi. Usilume chombo kitupu kwa kazi zingine zozote.

**ATHARI KWA MAZINGIRA:** SHIELD® 600 FS yaweza kudhuru samaki na wanyama. Kwa hivyo dawa yenyewe, mabaki yake au chombo kitupu hazifai kutumika au kutupwa kwa njia itakayochafua visima, mito, vidimbwi, au chemchemi za maji.

**HUDUMA YA KWANZA:**

**Ikitutwa kwa mapua:** Peleka mhadhiriwa palipo na hewa safi na umtuzi. Ikiwa hapumi, msaidie kupmua.

**Igusana na ngozi:** Safisha ngozi vizuri kwa maji yanayotirika kwa dakika 20.

**Ikingia kwa macho:** Fungua na kuyaosha macho kwa maji mengi kwa dakika 15.

**Ikimezwa:** Osha mdomo kwa maji. Usimfanye kutajika. Mtulize.

**PELEKA MDERUHI KWA DAKTARI**

**MAELEZO KUHUSU SUMU:** Yaweza kudhuru ikimezwa.

**Dalili za kusumika:** Huwasha kwenye macho, ngozi na mapua

**Makata:** Hakuna tiba maalum. Peleka mjeruhi kwa daktari.

**Arifa kwa daktari:** Tbu kulingana na dalili za kusumika

**Wakati wa kusumika, piga simu(masa 24) bila malipo kwa namba 0800720021/0800730030**

**ILANI KWA MTUMIAJI:** Dawa hii sharti itumie kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuanzima wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyosalama.

**THIBITISHO:** Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo lazima yafuatwe kikamilifu. Mhunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au maji na lazima akubali dawa katika hali hiyo.



**Greenlife**  
 CROP PROTECTION AFRICA

P. O. Box 24942, 00200 Nairobi, Kenya. Fax: +254 738 930257  
 Mbe: +254 735 54544, +254 722 553599, Tel: +254 20 24421012  
 info@greenlife.co.ke | www.greenlife.co.ke

Your Growth. Our Growth

Net Contents  
 (Kiasi) 1L