DESCRIPTION: SIGMA COMBI is a herbicide used to control grass weeds and broadleaves weeds in maize.

MODE OF ACTION: A selective systemic herbicide which acts as a photosynthesis inhibitor. It is a herbicide for the pre emergence and early post emergence control of annual grasses and selected broad leaved weeds in maize.

RECOMMENDATION FOR USE:

Crop	Weed	Application Rate	Remarks
Maize	Grasses and broadleaf weeds	3-4 ℓ per hectare in 150-250 ℓ per hectare or 250-400 mℓ in 16 ℓ of water knapsack	To be applied at pre-emergence stage of both crop and weeds. Apply immediately after planting onto fine, even and firm seed bed. Use high product rate if treatment is being done 7 days after rains.

Pre-Harvest Intervals: No harvest interval, **SIGMA COMBI** is applied in pre-emergence of crop and weeds.

Re-Entry Period: It is safe to enter sprayed area after 12 hours have passed.

HAZARD/ PRECAUTION: SIGMA COMBI has low mammalian toxicity and is classified as harmless to non-targeted organisms. **SIGMA COMBI** is highly toxic to aquatic organisms and is moderate to high persistence in the environment.

SAFETY PRECAUTIONS: Keep locked up and out of reach of children. Keep original container in a cool well ventilated place. Keep away from food, drinks and animal feeding stuffs. When using do not eat, drink or smoke. Do not breathe vapour. Avoid contact with skin and eyes. Do not empty into drains. Wear suitable protective clothing, gloves and face/eye protection. Use only in well ventilated areas. This material and its container should be disposed off as a hazard waste

STORAGE: Store in original tightly closed container in a cool dry place, safely locked, away from food, animals feedstuff, children and unauthorized persons. Advisable to keep temperature below 35°C.

DISPOSAL: Triple rinse or preferably pressure rinse container before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on-site. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, prefer the disposal on temperature incineration.

FIRST AID INSTRUCTIONS: Remove the affected person from danger zone to a well-ventilated room or to fresh air, and protect from chilling. Do not administer anything by oral route and do not try to make vomit, call a treatment centre for poisoning cases or a physician. Take the label where possible. **If inhaled:** Immediately remove to fresh air. Call a physician immediately. **If in contact with skin:** Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water. **If in contact with the eye:** Rinse immediately and thoroughly with plenty of water during at least 10 to 15 minutes. Eyelids should be held away from as is to ensure thorough rinsing. Seek medical advice if pain or redness persists. **If swallowed:** Call a physician immediately and show the label. Do not induce vomiting.

TOXICOLOGICAL CLASSIFICATION (WHO/FAO): II- Moderately Hazardous

SYMPTOMS OF POISONING: No adverse effect is likely as a result of normal handling and usage. Ingestion may produce nausea, vomiting and diarrhea.

NOTICE TO PHYSICIAN: Antidote: No specific antidote. **Treatment:** Treat symptomatically.

INCASE OF POISONING: Call the 24 hour Toll Free Numbers: **0800720021/0800730030**

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the pest control product act to use or store pest control product under unsafe conditions.

WARRANTY: Sellers guarantee is limited to the terms set out in the label. The buyer assumes the risk to the persons or property arising from the use of this product and accepts the product on that condition.

HERBICIDE

KIUA MAGUGU

SIGMA COMBI SC

A HERBICIDE FOR PRE-EMERGENCE AND EARLY POST EMERGENCE CONTROL OF ANNUAL BROADLEAF WEEDS AND GRASSES IN MAIZE.

NI KIUA MAGUGU CHA KUDHIBITI MAGUGU AINA YA NYASI NA YALE YALIYO NA MATAWI MAPANA KWENYE MIMEA YA MAHINDI

GUARANTEE (DHAMANA) ..... Acetochlor 250 g/ℓ, Atrazine 250 g/ℓ

REGISTRATION NUMBER (NAMBARI YA USAJILI) ..... PCPB (CR) 1008

COMMERCIAL AND AGRICULTURAL CLASS (KUNDI LA BIASHARA NA KILIMO)
READ THE LABEL BEFORE USING (SOMA KIBANDIKO KABLA YA KUTUMIA)
KEEP LOCKED UP OUT OF REACH OF CHILDREN (WEKA MBALI NA WATOTO)
IN CASE OF POISONING CALL TOLL FREE NO-24 HRS (WAKATI WA DHARURA PIGA SIMU BILA MALIPO KWA NAMBARI - MASAA 24)- 0800720021/0800730030

Registrant/ Msajiri:
WILLOWOOD Ltd Unit 2304, Wilborne Commercial Centre No 8 Java Road, HONG KONG

Manufacturer (Mtengenezaji):
Arysta LifeScience South Africa
7 Sunbury Office Park, Douglas Saunders Drive
La Lucia Ridge, 4019 SOUTH AFRICA | Tel: +27 31 514 5600

Distributor (Wakala / Msambazaji):
Arysta LifeScience (K) Ltd
Tulip House, 5th Floor Muindera Road
P.O. Box 30335 - 00100 G.P.O. Nairobi - KENYA
Tel: +(254) 720 671 282

Shelf life: Two years from the date of manufacture if stored in unopened original container in a dry cool place. **(Uasiha Rafuu:** Miaka miwili baada yatarehe ya kutengenezwa iwapo dawa iti kati fachwe kwa chombo cha ake asili bila kufunguliwa na wepesivumo baridi na jotoyenye)

Date of Manufacture (Tarehe ya kutengenezwa):
Batch N° (Nambari ya kifurusi):
Expiry Date (Tumia kabla ya):
} Seven packaging

Member of Agrochemical Association of Kenya AAK Reg No: 126

NET CONTENT (UJAZO)

1 ℓ

Arysta LifeScience

DRAFT LABEL APPROVED SUBJECT TO SUBMISSION OF FINAL LABEL AND PACKAGE VALID FOR 12 MONTHS ONLY
09 AUG 2018
PEST CONTROL PRODUCTS BOARD

MAELEZO KWA UJUMLA: SIGMA COMBI ni dawa ya kuangamiza kwenye ya aina ya nyasi magugu ya majani mapana kwa mimea ya mahindi. **SIGMA COMBI** huzuimbia kwenye kwenye namnagugu na kuzuia utengenezaji wa chakula. Ni dawa inayo uzuia kwenye ambazo hazija ota na ambazo zimeota zingali changa kwenye mimea ya mahindi

MAELEZO YA KIUFUNDI YA DAWA:

Mmea	Mimea ya Magugu	Kipimo cha dawa	Maelezo
Maize	Nyasi na magugu yaliyo na matawi mapana	Lita 3-4 kwenye hekta moja ndani ya lita 150-250 za maji au mililita 250-400 kwenye tumba la lita 16 za maji.	Nyunyizia kwenye udongo mwororo na chabiti mara moja baada ya kupanda na kabla ya mimea na kwekwe kuota. Tumia kiwango cha juu zaidi kama dawa inanyu - nyizwa baada ya siku saba baada ya mvua.

Muda salama wa kuvuna: Hakuna muda salama wa kuvuna kwa vile **SIGMA COMBI** hunyunyizwa kabla ya mimea na magugu kuota.

Muda salama wa kuingia shambani: Baada ya masaa 12.

TAHADHARI: SIGMA COMBI ina kiwango cha chini cha sumu kwa mammal an. Ni sumu kwawanyama wa majini na hudumu kwa muda kwenye udongo **Uhalabatanaji: SIGMA COMBI** ichanganyika vizuri na madawa ya kuwaida ya kuua wadudu na kuangamiza magonjwa. Muulima anaimizwa kufanya majaribio kabla ya kunyunyizia sehemu kubwa ya shamba.

USALAMA TAHADHARI: Weka mbali na watoto kwenye chumba kilichofungwa. Hifadhi kwenye chombo chake cha asili kwenye chumba chenye hewa safi. Weka mbali na chakula, maji ya kunywa na malisho ya wanyama. Wakati wa kupiga dawa shambani, usile chakula, usinywemaji, wala uvuta sigara. Epuka utusitasi wa dawa. Zita dawa isiguswe na ngozi au macho. Zata dawa isitiririke kwenye mfereji wa maji. Vaa nguo ili kujikinga wakatiwa kupiga dawa. Pia vaa glavu na mask ya macho. Fanya dawa ukiwa sehemu yenyehewa ya kutosha tu. Beada ya kutumia dawa pamoja na chupa yake iteketezwe katika maeneo maalum ya kuteketeza mabaki ya dawa. **Kama Kumeza: Usimtapishe** kutafta matibabu ushauri maramoja na kuonyesha kontena hii au studio.

KUTEKETEZA MABAKI YA DAWA WA VYOMBO TUPU: Kuendoksana ya bidhaa na kuduma yake ya ufungaji pamoja na kabla ya kuweji hika. Ustupe kontbi na mabowwa, mito, mitereji ama maamnde maisabe ya bidhaa. Osheni nyasi zilizochafuliwa na majina kwisha kuvusanya maji kwa matikabu. Sofisha pakiti ya dawa iliyoisha kwa kusuasa mara tatu utobos au uboryeze na ungue au unfermbali sa maji. Mchanganyiko wa dawa ambao haujatumika au maji yalio osha vyombo nilazima yatupiliwe mbali salama na ambapo pame idhinishwa.

MADHARA KW MAZINGIRA: SIGMA COMBI ni sumu kwa samaki, viumbe vingine vinavyoishi kwenye maji na nyuki. Inaweza kudhuru ndege na wanyama. Usitishe mazingira kwa kunyunyiza, kumwaga mabaki ya dawa kwasasa; mitaba wa dawa kwa mito, shimo au ardhinbwi na chemichemi. Usiny unyize cawa hii wakati nyuki wanatembelea maua. Usiruhusi mifugo na wanyama wa porikingia mabaki dawa hii mnyunyuziwa.

MAELEZO YA HUDUMA YA KWANZA: Ikipumuliwa: Toa mgonjwa mahali penye tatari na umpumzishe mahali ambapo nisalama. Kama mgonjwahwa ana shida ya kupumua, msaidie apumue, kasha umpeleke kwa daktari. **Ikimwagikia ngozi:** Vua mavazi ya hajilinga na osge kutumia maji safi na sabuni. Iwapo muumivu itaendelea, muone daktari. **Ikingia kwa macho:** Fungua macho asiribe kwa maji safi yanayotiririka kwa muda wa dakika 15. Iwapo maumivu yataendelea, muone daktari. **Ikimeza:** Usimfanye mgonjwa atapike. Mpeleke mekono, kasha umuone daktari.

MAELEZO KUHUSU SUMU: SIGMA COMBI ni sumu ikimezwa na ikingia kwa macho. Pia ni sumu ikipenya kwenye ngozi.Dalili za sumumka: Dalili za kusumika ni kama zile kuwashwa na ngozi. Kwanjo kidubwa cha sumuhu athiri 'Central Nervous System'. Tibu: Hakuna kifaa cha dare. Maelezo kwa Daktari: Toa mjamuzi wenye hatari na umpumzishe mahali pasipo na joto wala baridi. Mtibu mjeruhi kulingana na dalili za maumivu.

KWA HALI YA KUSUMIKA: Piga Namba ya simu **(MASAA 24) 0800 720 021/0800 730 030. ILANI KWA**

MTUMIAJI: Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye lebo hii. Nikatio chini ya sheria za madawa ya kuangamiza wadudu wahatibitu kutumia au kuhifadhi dawa hii katika hali isiyo salama.

THIBITISHO: Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo nilazima yafuatwe kikamilifu. Mnunuzihuvuma akubali hatari zozote zinazotokeza kutokana namatumizi ya dawa hii kwa biradamu au mali na kazime akubali dawa hii katika hali hiyo.