

MAELEZO KUHUSU DAWA

SUPREMO ina dawa aina ya alphacypermethrin na hutumika kudhibiti wadudu aina ya aphids, thrips, whiteflies na wengine kwenye maharagwe.

JINSI YA KUTUMIA

Mmea	Wadudu	Kipimo
Maharagwe	Aphids	mililita 10 kwa lita 20 za maji
"	Thrips	"
"	Whiteflies	"

Wakati wa kungoja kabla kuvuna: siku saba

wakati wa kungoja kabla kurudi shambani: masaa 12

Muda wa kungoja kabla kunyunyizia dawa tena: Siku 14

TAHADHARI

Chunga isiingie machoni, kuguza ngozi, kuvuta mvuke au kuimeza. Usivute sigara au kula wakati unaposhugulikia dawa hii. Wakati unaposhugulikia dawa hii vaa mavai ya kujikinga kama vile mifuko ya mikono na miwani ya kuzuia macho. Oga vizuri na ufue nguo hizo za kazi kisha uvae nguo zilizo safi.

UHIFADHI

Hifadhi dawa ndani ya chombo chake asili na ufunge vizuri kwenye mahali penye giza na mbali na watoto, mifugo na pia vyakula.

usiweke dawa karibu na moto au vyombo vyenye kutoa joto.

UTUPAJI

Usitumie vyombo vilivyo vitupu kwa njia yeyote. Suza mara tatu na uvitoboe halafu utupe kulingana na sheria na kanuni za utupaji za nchi.

Iwapo dawa imemwagika, zoa mchanga mkavu au (vumbi la mbao) kisha uinyunyizie na uzike.

Ikiwa kuna haja ya kutupa kiasi kikubwa cha Supremo, pata mashauri kutoka kwa watengenezaji au wale ambao wanahusika na ugawaji dawa hii.

Iwapo ni kiasi kidogo cha dawa punguza nguvu na maji na kisha unyunyizie kwenye sehemu ambazo hazina mimea yenye mazao.

HATARI KWA MAZINGARA

SUPREMO ni hatari kwa samaki na viumbe wengine wa majini. Usitupe dawa au chombo kwenye maji baada ya kushugulikia Bestox.

HUDUMA YA KWANZA

Inapovutwa: mpeleke mhusika katika sehemu yenye hewa safi. Mpeleke mjeruhi kwa daktari.

Inapoingia machoni: Osha kwa maji mengi kwa muda wa dakika 15.

Inaposhika ngozi: Toa nguo zilizomwagikiwa na uoshe kwa maji ya sabuni.

Inapomezwa: Usijaribu kujitapisha. Mpeleke mjeruhi kwa daktari.

MAARIFA JUU YA SUMU

Dalili ya sumu: Unyonge, kuishiwa na nguvu na kuumwa na kichwa.

Mwongozo kwa daktari

Supremo ina mchanganyiko wa madawa hatari ambayo yanaweza kusababisha kuharibika kwa mapafu, ikipenya wajkati wa kutapika. Wakati mhusika anapotapika, tumia mrija maalum wakitibabu. Hakuna kinga ya madhara kama haya. Shughulikia mgonjwa kulingana na dalili alizo nazo.

Hakuna dawa ya kuharibu sumu ya dawa hii,

Wakati wa kusumika piga simu bila malipo kwa Nambari: 0800730030 ama 0800720021 (MASAA ISHIRINI NA MANNE)

ILANI KWA MTUMIAJI

Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyo salama.

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokea kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.

DESCRIPTION

SUPREMO contains alphacypermethrin, and is used to control thrips, aphids and whiteflies on french beans.

MODE OF ACTION

Alphacypermethrin acts by contact and stomach action on the insects and affects the nervous system.

DIRECTIONS FOR USE

Supremo is used as a foliar spray.

CROP	PEST	RATE	PHI
Frenchbeans	Aphids	10mls/20L water	7days
"	Thrips	"	"
"	Whiteflies	"	"

Application Interval = 14 days

PHI = 7 days

Re-entry period = 12 Hours

HAZARDS / PRECAUTIONS

Avoid inhalation, contact with skin or eyes and ingestion. Do not eat or smoke when applying the product. When handling concentrate wear protective clothing including gloves and goggles. Avoid breathing spray mist and do not spray into the wind. After spraying, bathe, wash work clothes and change into clean clothes.

STORAGE

Store in tightly closed original containers in a cool dark place away from children, livestock and food. Do not store near sources of heat or fire.

DISPOSAL

Puncture holes to the empty containers and dispose in accordance to the national laws and regulations on disposal.

To dispose large quantities of Supremo contact manufacturer/supplier. Small quantities should be diluted and sprayed on non crop areas. Do not re-use container for any other purpose.

Do not pour unused Supremo or sprayer washings into waterways.

If concentrate spills absorb spillage with dry soil or sawdust and collect and bury.

Clean contaminated area thoroughly with water and drain away from water sources.

ENVIRONMENTAL HAZARD

SUPREMO is toxic to fish and other aquatic organisms. Do not contaminate water bodies with re-usage of containers.

FIRST AID INSTRUCTIONS

If inhaled: Remove the person to fresh air. Call a doctor

If in eyes: Flush with plenty of water for at least 15 minutes.

If on skin: Remove contaminated clothes and wash with water and soap.

If swallowed: DO NOT induce vomiting, call a doctor.

TOXICOLOGICAL INFORMATION

Symptoms of poisoning

Headache, fatigue and anorexia

Antidote: None. Treat symptomatically.

Note to the doctor

Aromatic hydrocarbons can cause lung damage if aspirated during vomiting. Induce vomiting with an endotracheal tube in place. There is no Antidote. Treat symptomatically.

In case of poisoning call Toll Free Number: 0800730030 or 0800720021(24HRS)

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under The Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY

Sellers guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to the persons or property arising from the use or handling of this product and accepts the product on that condition.

SUPREMO[®] 100 EC

INSECTICIDE/(KIUA DUDU)

For the control of thrips, aphids and whiteflies, on frenchbeans
(Kiua dudu cha kudhibiti wadudu kama thrips, aphids na whiteflies
kwenye mishiri.)

GUARANTEE/(DHAMANA):
Alphacypermethrin

Contains
10% W/V

**COMMERCIAL AND AGRICULTURAL CLASS
(KUNDI LA BIASHARA NA KILIMO)
REGISTRATION NO(NAMBARI YA USAJILI). PCPB (CR) 1280**

**KEEP LOCKED AWAY
FROM CHILDREN**

**(WEKA IMEFUNGIWA
MBALI NA WATOTO)**

**READ THE LABEL CAREFULLY BEFORE USE AND USE ONLY AS DIRECTED
(SOMA MAELEZO KWA MAKINI KABLA YA KUTUMIA)**



CAUTION
May cause harm

(TAHADHARI
Inaweza kudhuru)



Flammable
Inaweza
Kupata
Moto

**IN CASE OF POISONING CALL TOLL FREE NUMBER:
(WAKATI WA KUSUMIKA PIGA SIMU BILA MALIPO KWA NAMBARI:
0800730030 or/au 0800720021(24HRS)**

Shelf Life: Two (2) years in original
unopened container after date of
manufacture stored in a cool dry place

(Maisha Rafuni: Miaka 2 baada ya tarehe ya
kutengenezwa kwenye chombo chake asili
kilichofungwa mahali pakavu pasipo na joto
jingi.)

Date of Manufacture/(Tarehe ilipotengenezwa):

Expiry Date(Tumia Kabla ya):

Batch No./(Nambari ya furushi):

NET CONTENTS

1Litre

(UZITO KAMILI)

Agent/Ajenti:

JUANCO SPS LTD.

Juanco Centre, Ngong Rd., Ngong Hills

P.O.Box 381, Karen 00502, Nairobi.

Tel: +254 (20) 2088754, 2088755

2088789, 2088754, 2048774, 3002398

Fax: +254 (20) 2088793

Mobiles: +254 (722) 207805, 207806

Email: info@juancogroup.com

Website: www.juancogroup.com

**DRAFT LABEL APPROVED SUBJECT TO
SUBMISSION OF FINAL LABEL AND PACKAGE
VALID FOR 3 MONTHS ONLY**

06 FEB 2019

PEST CONTROL PRODUCTS BOARD

Member of
Agrochemical
Association
of Kenya
A.A.K No 273



Manufactured by/(imetengenezwa na): Sinochem Ningbo
Chemicals Co. Ltd. 21 Jinagxia Street, Ningbo, China,
Xiepu Town, Zhenhai District, Ningbo P.R. China



**HARMFUL
INADHURU**