

Talstar® 10 WP

DIRECTIONS FOR USE:

Talstar® 10WP is a residual insecticide for control of Mosquitoes in IRS programmes. All support surfaces can be treated such as bricks, wood, mud, concrete, galvanized iron, cement and others. Dissolve the contents of the sachet (62.5g) in 10 litres of water and then stir. Spray the surfaces to wet (40 ml/m²). Talstar has a residual protection period exceeding 6 months depending on treated surface material.

SAFETY DIRECTIONS

Store in a tightly closed container away from reach of children, food stuff and fire. Keep away all pets and food materials from the area to be sprayed. Wear gloves, masks, long-sleeved shirts, trousers and gumboots before opening the sachet, mixing and spraying the product. Do not eat, drink or smoke while handling the product. Before eating, drinking or smoking after working, wash hands and any other parts of contaminated skin thoroughly with plenty of water and soap. After each usage, clean with plenty of water all tools, equipment and all protective clothing. Open all doors and windows for good ventilation and clean the floor after treatment.

RE-ENTRY

Re-entry into treated house is safe 1 hour after treatment.

WHO Classification

Talstar 10WP belongs to class II, Moderately hazardous, HARMFUL.

STORAGE

Store in cool dry place. Do not change storage containers. Destroy empty containers. Do not re-use empty containers. Do not contaminate other pesticides, fertilizers, water and feed by storage.

DISPOSAL

Do not contaminate ponds, water or sewage systems through disposal of washing from cleaning equipment.

Collect sachets and dispose them off as recommended

away from water sources or settlements in accordance with national legislation and regulations.

FIRST AID INSTRUCTIONS

If inhaled: Move the person to fresh air. Call a doctor.

If in eyes: Flush eyes with plenty of water for at least 15 minutes.

If on skin: Remove contaminated clothes and wash skin with plenty of water and soap.

If swallowed: Rinse mouth with clean water. Dilute by giving 1-2 glasses of water. Do not induce vomiting. Obtain immediate medical attention.

PRECAUTIONS

Avoid contact with skin and eyes. Do not eat, drink, or smoke when applying the product. Bifenthrin is highly toxic to fish and aquatic organisms. Keep out of drains and water courses.

TOXICOLOGICAL INFORMATION

-Symptoms of poisoning: Tremor, convulsions and bleeding from the nose. If one of these symptoms manifests, stop working and perform First Aid immediately and then see a doctor.

-Antidote: Has no antidote.

-Note to the physician

Bifenthrin is a synthetic pyrethroid containing aromatic solvents.

If swallowed perform gastric lavage taking care to prevent aspiration of the gastric contents. Thereafter treat symptomatically.

IN CASE OF POISONING CALL TOLL FREE NUMBER:
0800730030 or 0800720021(24HRS)

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act, to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling this product and accepts the product on that account.

JINSI YA KUTUMIA

Talstar® 10 WP ni dawa ya kunyunyizia ndani ya nyumba ili kukabiliana na mbu wanaosambaza ugonjwa wa Malaria. Kuta au vieso za aina yote kama vile matope, mabati, mawe, mbao nyasi, simiti na kathalka zaweza kunyunyiziwa na dawa hii ili kuthibiti mbu wasambazao ugonjwa wa Malaria.

-Dawa hii ni lazima kuchanganywa na maji kabla ya kunyunyiziwa

Kutumia bomba la mkononi aina ya knapsack.

-Tia maji kufikia nusu ya tanki la bomba na weka pakiti yote gramu 62.5 kwenye chombo cha kunyunyizia halafu ongeza maji hadi ifike lita 10.

-Tikisa vizuri kuhakikisha dawa imepata kuchanganyika vyema na maji.

-Nyunyizia kwenye kuta hadi zilowe (mililita 40 kwa mita moja mraba).

Yaweza kulinda kwa zaidi ya miezi sita.

-Muda salama wa kuingia pahali paliponyunyiziwa dawa ni baada ya isaa limoja.

TAHADHARI

Hifadhi dawa hii mahali penye usalama mbali na watoto na moto.

- Waondoe wanyama wa nyumbani na aina yote ya vyakua kutoka sehemu za kunyunyizia dawa.

-Valia glavu, vikinga uso, shati la mikononi mirefu, longi na buti kabla ya kufungua na kunyunyizia dawa. Usile, kunywa, au kuvuta sigara unapotumia dawa.

-Baada ya kazi, osha mikono na sehemu zote ambazo zimepata dawa na maji mengi na sabuni kabla ya kula, kunywa au kuvuta sigara.

Kila baada ya kutumia, osha nguo na vyombo vyote ukitumia maji mengi masafi.

-Baada ya kunyunyizia dawa, fungua mlango na madirisha yote ya nyumba ili kupokea hewa safi.

UWEKAJI

Weka kwenye sehemu kavu isiyokuwa na joto. Usibadilishe vyombo inamowekwa dawa hii. Haribu kabisa vyombo vya dawa vilivyotumika na utupe kulingana na sheria za utupaji za nchi. Usitumie vyombo hivi tens.

UTUPAJI

-Usichafue vidimbwi na mitaro ya maji machafu kwa kuelekeza maosho yanayotokana na usafishaji wa vyombo vilivyotumiwa kwa unyunyiziwa.

MAAGIZO KWA DAKTARI

Hii dawa ni sumu wastani ikifutwa au kumezwa. Husababisha Mwasho ikiingia machoni lakini haisababishi mwasho kwenye ngozi. Usimpe aliyeathiriwa maziwa au chakula chochote. Tibu ili kupunguza wingi wa sumu mwilini.

DALILI ZA KUATHIRIWA

Kuteleleka, kuzimia na kutoka damu pua. Ukiona kati ya hizi dalili pokea huduma ya kwanza na umuone daktari haraka iwezekanavyo.

HUDUMA YA KWANZA

Inapovutwa: mpeleke mhusika katika sehemu yenye hewasafi. Muite daktari

Inapoiingia machoni: Osha kwa maji mengi kwa muda wa dakika 15

Inaposhika ngozi: Toa nguo zilizopata dawa na uoshe ngozi kwa maji na sabuni

Inapomezwa: Usimtapishe mhusika. Muite daktari mara moja.

WAKATI WA KUSUMIKA PIGA SIMU BILA MALIPO KWA NAMBARI: 0800730030 au 0800720021(24HRS)

ONYO

Ni sumu ikimezwa, vutwa au kuingia mwilini kupitia ngozi au macho. Huweza kuthuru nyuki, samaki, wanyama wa nyumbani, mifugo na hata wanyama na ndege wa porini.

ILANI KWA MTUMIAJI

Dawa hii sharti itumiwe kulingana na maagizo yalio kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyosalama

THIBITISHO

Dhamana ya muuzaji ni maelezo yalio kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnuuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima akubali dawa hii katika hali hiyo.

